

## STUDIES ON THE EFFECT OF COMBINATION OF BIOREGULATORS AND GROWTH REGULATORS ON FLOWERING AND YIELD OF MANGO

(*Mangifera indica* L) cv. BANGANPALLI

DHEERAJ, G<sup>1</sup>, BHAGWAN, A<sup>2</sup>, RAJ KUMAR, M<sup>3</sup> & SRIDHAR, D<sup>4</sup>

<sup>1,4</sup>College of Horticulture, Rajendranagar, Hyderabad, Telangana, India

<sup>2,3</sup>Fruit Research Station, Sangareddy, Medak, Telangana, India

### ABSTRACT

An experiment on the effect of bioregulators and growth regulators zinc sulphate @ 0.1%, paclobutrazol (PBZ) @ 3ml.m<sup>-1</sup> canopy diameter, Naphthalein acetic acid (NAA) @ 80 ppm, gibberilic acid (GA<sub>3</sub>) @ 20 ppm, 1-2-chloro-4pyridyl-3-phenyl urea (CPPU) @ 10ppm, potassium sulphate (K<sub>2</sub>SO<sub>4</sub>) @ 1% , their combinations utilizing BBCH scale of mango was conducted on flowering, fruit set and yield of mango cv. Banganpalli. Paclobutrazol alone or in combinations with other bioregulators has significantly increased the percent of flowering, panicle length and breadth when compared to control. Significantly highest number of fruits.panicle<sup>1</sup>, fruits.tree<sup>1</sup> and yield (23.7 % over control) were recorded in paclobutrazol alone followed by Naphthalein acetic acid (20.96% increases in yield over control).

**KEYWORDS:** Bioregulators and Growth Regulators, Fruit Set and Yield of Mango

**Received:** Mar 20, 2016; **Accepted:** Apr 07, 2016; **Published:** Apr 12, 2016; **Paper Id.:** IJASRJUN20163

### INTRODUCTION

Mango (*Mangifera indica* L.) is the premier fruit among the tropical fruits. India is the largest producer of mango in the world. The fruits of mango are valued because of its excellent flavor, appealing aroma, delicious taste, attractive shades of colour and nutritive value, which has attracted the world market. . In Telangana, mango occupies an area of 2.10 lakh ha, with a production of 1,894.9 M.T with a productivity of 9.0 T. ha<sup>-1</sup> (Anon, 2013). Although, India is the leading mango producing country, the development of mango orchards as an established industry has remained a distant goal and per hectare yields (2.45 t ha<sup>-1</sup>) are low in spite of great potential (Anon, 2013). There are number of reasons for the poor productivity, of which alternate or biennial bearing habit of most of the choicest commercial varieties of India is one of the important factors. The flowering phenomenon in mango appears to be a complex one. Research workers engaged in the field have not appreciably established the critical factors responsible for this phenomenon. Recommendations of horticultural scientists have not paved the way for a satisfactory solution to this problem. However, on account of continued research, this problem is better understood now than before. Since, most of the earlier studies were aimed at developing agro techniques to promote regular cropping, enough knowledge on the physiology of flowering is lacking. However, the studies of Davenport (2007), on the evidence of involvement of leaf generated floral stimulus have opened up new vistas in the direction of research on physiology of flowering in mango.

Various chemicals and growth regulators application have been standardised for enhancing vegetative

growth and uniform flowering in mango. Application of Zinc Sulphate ( $\text{ZnSO}_4$ ) @ 0.4% promoted vegetative growth as indicated by plant height, girth, and spread of young plants (Banik and Sen 1997). Application of Gibberellic acid ( $\text{GA}_3$ ) increased vegetative growth, number of secondary branches (Singh and Rajput 1990). Paclobutrazol (PBZ) @  $3\text{ml.m}^{-1}$  of canopy diameter at 120 days before bud break was found to improve flowering and yield in mango cv. Banganpalli (Bhagwan *et al.*, 2011). Naphthalein Acetic Acid (NAA) @ 80 ppm spray at 30 days before flowering was found to improve flowering in mango (Davenport, 2007). Application of CPPU (10 ppm) gave the best result in terms of increasing fruit retention, number of fruits per cluster and per plant, weight of one fruit, fruit volume and leaf area in mango cv. Arumanis (Notomedijo 1999).

Knowledge of the periodic biological events of a particular crop – bud breaks, flushing, flowering, and fruit development is an important tool for its agronomical management in relation with climate change (Hernandez Delgado *et al.*, 2011). Bleiholder *et al.* 1989 developed a two-digit decimal coding system for angiosperms, the BBCH-scale (Biologische Bundesantalt, Bundessortenamt und Chemische Industrie). Later three digits “extended BBCH-scale” (Hack *et al.*, 1992) was proposed for certain crops like grains, rape and sunflower, vegetables, pome and stone fruits, citrus, grape, strawberry, pomegranate, coffee, olive, Musaceae, persimmon, cherimoya. Keeping above fact in view, the phenological studies were conducted to assess BBCH- scale for mango cultivars like Totapuri (Shailender rajan *et al.*, 2011). The BBCH scale distinctly separates the various vegetative flushes occurring in the mango, as well as the terminal and axillary flowerings. All of which is important for the correct timing of general orchard management, particularly for disease and pest management, physiological disorders and weed control, flowering inhibition and effectiveness of fertilizers and PGRs application (Hernandez Delgado *et al.*, 2011).

## MATERIALS AND METHODS

The experiment was conducted on ten years old well grown, uniform statured trees of mango cv. Banganpalli. Trees are spaced with 8m and planted in square system. The statistical design adopted was Randomised block design with 8 treatments which were replicated thrice. Paclobutrazol concentration was calculated based on the diameter of the tree, and applied @ 3 ml per metre of canopy diameter (Bhagwan *et al.*, 2011). The required paclobutrazol was dissolved in 10 litre of water. Eighty milligrams of Naphthalein acetic acid (NAA) was dissolved in 50 ml of ethanol and diluted it in 1 litre of water to get 80 ppm of NAA. Ten gm of potassium sulphate ( $\text{K}_2\text{SO}_4$ ) was dissolved in 1 litre of water to get 1% of  $\text{K}_2\text{SO}_4$ . Twenty mg of (1-2-chloro-4pyridyl-3-phenyl urea) CPPU was dissolved in 50 ml of ethanol and diluted it in 1 litre of water to get 20 ppm of CPPU. Twenty mg of gibberellic acid ( $\text{GA}_3$ ) was dissolved in 50 ml of ethanol and diluted in 1 litre of water to get 20 ppm of  $\text{GA}_3$ . 0.1 gram of zinc sulphate ( $\text{ZnSO}_4$ ) was dissolved in 1 litres of water to get 0.1% of  $\text{ZnSO}_4$ . Data on percent flowering, panicle length and breadth, fruitset. panicle<sup>1</sup>, fruits. Tree<sup>1</sup> and yield were recorded. Twenty shoots were randomly tagged (from North, South, East and West directions) and no of tagged shoots which had flowered was recorded and expressed as percentage of flowering. The panicle length and breadth, fruit set of ten randomly selected shoots were recorded the mean was calculated. The total number of fruits harvested.tree<sup>-1</sup> was counted after harvest and expressed as number of fruits.tree<sup>-1</sup>. The average fruit weight was computed by dividing the total yield (kg per tree) and number of fruits per tree of the respective treatment.

## RESULTS AND DISCUSSIONS

### Flowering Parameters

The results on percent flowering (%) of mango cv. Banganpalli sprayed with different combinations of

bioregulators and growth regulator sprays are presented in the table 1.

There was significant difference among combination of bioregulators and growth regulator sprays application with respect to percent flowering (%). Trees sprayed with paclobutrazol (91.67) has significantly increased the percent flowering which was on par with application of Naphthalein acetic acid (73.33) and gibberillic acid (75). Minimum flowering percent was recorded in untreated control (35), followed by application of potassium sulphate + (1-2-chloro-4pyridyl-3-phenyl urea) (45) which were on par with application of (1-2-chloro-4pyridyl-3-phenyl urea) (61.67), zinc sulphate + paclobutrazol (56.66), zinc sulphate + (1-2-chloro-4pyridyl-3-phenyl urea) (55), potassium sulphate + gibberillic acid (51.66) and potassium sulphate + Naphthalein Acetic Acid (50).

Paclobutrazol application and Naphthalein acetic acid spray has significantly increased the per cent flowering compare to control (Table 1). However, the paclobutrazol was significantly more effective in increasing the flowering percentage. Gibberellins, a group of plant growth hormones were reported to be inhibitory to flowering (Kachru *et al.*, 1971), in mango and the available evidence suggests the flower promotive effect of paclobutrazol in mango due to its anti - gibberellin activity (Quinlan and Richardson, 1984). Hence, in the present investigation the increase in the percent (%) flowering of mango by paclobutrazol was due to its anti - gibberellin activity. Increase in per cent flowering (%) by application of Paclobutrazol was earlier reported by Rao *et al.* (1997) in mango cv. Alphonso, Shinde *et al.* (2001) also reported similar increase in per cent flowering (%) by application of Paclobutrazol in mango cv. Alphonso. Zora Singh *et al.* (2000) also reported similar increase in per cent flowering (%) by application of Paclobutrazol in mango. Burondkar *et al.* (2000) reported similar increase in per cent flowering (%) when the trees were sprayed with paclobutrazol in mango cv. Alphonso. Vijayalaxmi and Srinivasan (2000) also reported similar increase in per cent flowering (%) by application of Paclobutrazol in mango cv. Alphonso. Similar increase in per cent flowering (%) by application of Paclobutrazol was obtained by Kumar Raj *et al.* (2005) in mango cv. Banganpalli and by Orwintinee *et al.* (2008) in mango cv. Irwin. The naphthalein acetic acid spray has also significantly increased the flowering in mango cv. Banganpalli in the present investigation when compared with the control. Increase in flowering with spraying of Naphthalein acetic acid was reported in mango cv. Succary Abiad (Wahdan *et al.*, 2011). Naphthalein acetic acid which is considered as flowering hormones in some crops might have increased the latent flowering factors in the mango and resulted in overall increase in flowering in mango cv. Banganpalli when compared to control in the present investigation.

The results on panicle length after application of different combination of bioregulators and growth regulator sprays are presented in the table 2. There is significant difference among different combination of bioregulators and growth regulator sprays with respect to panicle length. Minimum panicle length was recorded in untreated control (19.30), followed by application of (1-2-chloro-4pyridyl-3-phenyl urea) (20.88), gibberellic acid (21.81), zinc Sulphate + gibberellic acid (24.32), potassium sulphate + gibberellic acid (24.34) and zinc sulphate + naphthalein acetic acid (24.97). Trees sprayed with paclobutrazol (30.57) has significantly increased the panicle length compared to control (19.30).

Paclobutrazol has significantly increased the panicle length compare to control (Table 2). Similar increase in panicle length with the application of paclobutrazol in mango was reported by Desai and Chundawat (1994) in mango. Vijayalaxmi and Srinivasan (1998) also earlier reported increase in panicle length in mango cv. Alphonso when trees treated with paclobutrazol @ 10 ml.tree<sup>-1</sup>.

The results on panicle breadth after application of different combination of bioregulators and growth regulator

sprays are presented in the table 3. There is significant difference among different combination of bioregulators and growth regulator sprays with respect to panicle breadth. Trees sprayed with paclobutrazol (22.2) has significantly increased panicle breadth which was on par with application of zinc sulphate + (1-2-chloro-4pyridyl-3-phenyl urea) (21.82) and zinc sulphate + paclobutrazol (18.51) compared to control (8.33).

Paclobutrazol could able to increase the panicle breadth compare to control and naphthalein acetic acid (Table 3). However, Winston (1992) in mango cv. Kensington and Orwintinee *et al.* (2008) in mango cv. Irwin reported that the panicles of paclobutrazol treated trees were considerably shorter than those of control trees. The discrepancy in the finding of present investigation to the earlier reports regarding panicle may be due to varietal change, time of applications and dosage of paclobutrazol. However, increase in panicle length and breadth of paclobutrazol treated trees might be beneficial for increase the number of hermaphrodite flowers per panicle. This may cause for better fruit set over the control.

### Fruit Set Parameters

The results on fruit set per panicle after application of different combination of bioregulators and growth regulator sprays are presented in the table 4. The data revealed that there is significant difference among different combination of bioregulators and growth regulator sprays with respect to fruit set per panicle of mango. Trees sprayed with paclobutrazol (13.13), zinc sulphate +(1-2-chloro-4pyridyl-3-phenyl urea) (13) have significantly increased fruit set per panicle which were on par with the application of potassium sulphate+ paclobutrazol(12.80), potassium sulphate + naphthalein acetic acid(11.33), potassium sulphate + gibberellic acid (10.86), zinc sulphate + gibberellic acid (10.13). However, zinc sulphate+ naphthalein acetic acid (8.2), potassium sulphate+(1-2-chloro-4pyridyl-3-phenylurea) (8.33), zinc sulphate + paclobutrazol (8.43) and gibberellic acid (8.86) could not significantly increase the fruit set per panicle.

The prolonged flowering period (Days taken for full flower opening from panicle initiation) might have caused for better fruit set in paclobutrazol treated trees compare to control.

Application of paclobutrazol and zinc sulphate + (1-2-chloro-4pyridyl-3-phenyl urea) increased fruit set panicle<sup>-1</sup> compared to control and naphthalein acetic acid (Table 4). Similar increase in fruit set panicle<sup>-1</sup> in response to paclobutrazol application was recorded by GoGuey (1990) in cv. Valencin. Increase in fruit set panicle<sup>-1</sup> in response to paclobutrazol application was earlier reported by Iyer and Kurian (1992) in mango cv. Alphonso. Zora Singh *et al.* (2000) also reported increase in fruit set panicle<sup>-1</sup>, when trees were treated with paclobutrazol in mango cv. Dashehari. Increase in fruit set panicle<sup>-1</sup> in response to paclobutrazol application was earlier obtained by Cardenas and Rojas (2003) in mango cv. Tommy Atkins. Kumar Raj *et al.* (2005) also reported increase in fruit set panicle<sup>-1</sup>, when the trees were treated with paclobutrazol in cv. Banganpalli. Increased fruit set panicle<sup>-1</sup> in response to the application of paclobutrazol was earlier reported by Orwintinee *et al.* (2008) in mango cv. Irwin.

Increase in fruit set panicle<sup>-1</sup> in response to application of zinc sulphate + (1-2-chloro-4pyridyl-3-phenyl urea) might be due to the increase in fruit set after the application of synthetic cytokinin (1-2-chloro-4pyridyl-3-phenyl urea).

Similar increase in fruit set panicle<sup>-1</sup> in response to application of (1-2-chloro-4pyridyl-3-phenyl urea) was recorded by Notodemedjo (2000) in mango cv. Arumanis and Burondkar *et al.* (2009) in mango cv. Alphonso. Increase in fruit set panicle<sup>-1</sup> in response to application of zinc sulphate, was reported by Negi *et al.*, (2011) in mango cv. Dashehari.

## **Yield Parameters**

The results on total number of fruits produced on a tree after application of different combination of bioregulators and growth regulator sprays are presented in the table 5.

The data revealed that there is significant difference among combination of bioregulators and growth regulator sprays with respect to number of fruits per tree of mango. Trees sprayed with application of paclobutrazol (133) has significantly increased number of fruits per tree which was on par with application of (1-2-chloro-4pyridyl-3-phenylurea) (131.66), zinc sulphate + paclobutrazol (131.33) and potassium sulphate+ paclobutrazol (130.00). Minimum number of fruits per tree was recorded in untreated control (110.66), zinc sulphate + gibberillic acid (112.33) and gibberillic acid (113).

Trees sprayed with paclobutrazol have recorded increase in per cent flowering (Table 1), prolonged days taken for 50% flowering and days taken for full flower opening which inturn increased the fruit set per panicle (Table 4). This increase in fruit set per panicle was responsible for the increase in fruit number in the paclobutrazol sprayed trees in the present investigation. The correlation between intensity of flowering, perfect flowers and better fruit set and subsequent increase in total number of fruits per tree and yield was earlier reported by Burondkar and Gunjate, (1993) in mango. Paclobutrazol application has significantly increased the number of fruits per tree compare to control and naphthalein acetic acid spray (Table 5). Similar increase in number of fruits tree<sup>-1</sup> was earlier reported by Kumar Raj *et al.* (2005) in mango cv. Banganpalli. Increase in number of fruits in response to paclobutrazol application was earlier obtained by Cardenas and Rojas (2003) in mango cv. Tommy Atkins and by Orwintinee *et al.* (2008) in mango cv. Irwin. Kumbhar *et al.* (2007) also reported similar increase in number of fruits in mango cv. Kesar.

The results on fruit weight (gm) of mango cv. Banganpalli sprayed with different combination of bioregulators and growth regulator sprays are presented in the table 6. There was significant difference among different combination of bioregulators and growth regulator sprays application with respect to fruit weight (gm). Trees sprayed with application of naphthalein acetic acid (315.67) has significantly increased the individual fruit weight which was on par with application of paclobutrazol (309.33), zinc sulphate + paclobutrazol (309) and potassium sulphate+ naphthalene acetic acid (307.33) compared to untreated control (296.67).

There was significant difference among different combination of bioregulators and growth regulator sprays with respect to fruit weight (gm) (Table 6). Paclobutrazol and NAA were both equally increased the fruit weight. Similar increase in fruit weight was earlier reported by Nav Prem Singh *et al.* (2002) in mango cv. Dasher treated with 100 ppm of NAA and Wahdan *et al.* (2011) in mango cv. Succary Abiad treated with NAA @ 60 ppm. Paclobutrazol also increased the fruit weight in mango cv. Baneshan (Kumar Raj *et al.* 2005). Increase in fruit weight was earlier reported by Razzaq *et al.* (2013) in kinnow mandarin when the trees were treated with 0.6% zinc sulphate. Similar increase in fruit weight was earlier reported by Burondkar *et al.* (2009) with the application of Potassium sulphate in mango cv. Alphonso.

The results on yield per tree after application of different combination of bioregulators and growth regulator sprays are presented in the table 7. The data revealed that there is significant difference in yield (kg.tree<sup>-1</sup>) among different combination of bioregulators and growth regulator sprays. Trees sprayed with application of paclobutrazol (40.6) have significantly increased the yield per tree which was on par with application of naphthalein acetic acid (39.7) and zinc sulphate + paclobutrazol (39.61). Minimum yield was recorded in untreated control (32.82).

Paclobutrazol and NAA were both equally increased the yield. The increase in intensity of flowering, better fruit set and fruit weight in paclobutrazol treated trees have ultimately increased the yield of mango (23.7% over control) in the present investigation. Burondkar and Gunjate (1993) also found similar correlation between flowering, fruit set, fruit weight and yield of mango in response to paclobutrazol application. Similar increase in yield of mango in response to paclobutrazol application was obtained by various workers like Singh and Dhillon (1992) in cv. Kensington pride, Zora Singh and Dhillon (1992) also reported similar increase in yield in cv. Dashehari. Increase in yield of mango in response to paclobutrazol application was obtained by Iyer and Kurian (1992), Ram and Tripathi (1993) in cv. Dashehari. Desai and Chundawat (1994) also reported similar increase in yield in mango cv. Alphonso. Patil and Talathi (1999), Shinde *et al.* (2002) earlier reported similar increase in yield in cv. Alphonso. Increase in yield of mango in response to paclobutrazol application was earlier reported by Kumar Raj *et al.* (2005) in mango cv. Banganpalli. Orwintinee *et al.* (2008) also obtained similar increase in yield in cv. Irwin and Kumbhar *et al.* (2007) reported similar increase in yield in response to paclobutrazol application in mango cv. Kesar.

Similar increase in yield in response to the application of zinc sulphate was reported by Negi *et al.* (2011) in mango cv. Dashehari, Singh *et al.* (2009) in mango, Lolaei *et al.* (2012) in strawberry. Abdollahi *et al.* (2010) also earlier reported similar increase in yield in response to the application of zinc sulphate in Strawberry cv. Selva.

Rawash *et al.* (1998a) in mango and Shinde *et al.* (2002) also earlier reported similar increase in yield in response to the application of naphthalein acetic acid in mango cv. Alphonso, Application of naphthalein acetic acid increased yield in mango cv. Dashehari (Singh *et al.*., 2005).

**Table 1: Effect of Combination of Different Bioregulators and Growth Regulator Sprays on Percent Flowering of Mango Cv. Banganpalli**

Treatment	Mean
T <sub>1</sub> : Gibberilic acid	75.00 <sup>ab</sup>
T <sub>2</sub> : Naphthalein Acetic Acid	73.33 <sup>ab</sup>
T <sub>3</sub> : Paclobutrazol	91.67 <sup>a</sup>
T <sub>4</sub> : (1-2-chloro-4pyridyl-3-phenyl urea)	61.67 <sup>bc</sup>
T <sub>5</sub> : Zinc Sulphate+Gibberilic acid	48.33 <sup>bc</sup>
T <sub>6</sub> : Zinc Sulphate+Naphthalein Acetic Acid	35.00 <sup>c</sup>
T <sub>7</sub> : Zinc Sulphate+Paclobutrazol	56.67 <sup>bc</sup>
T <sub>8</sub> : Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	55.00 <sup>bc</sup>
T <sub>9</sub> : Potassium Sulphate +Gibberilic acid	51.67 <sup>bc</sup>
T <sub>10</sub> : Potassium Sulphate +Naphthalein Acetic Acid	50.00 <sup>bc</sup>
T <sub>11</sub> : Potassium Sulphate +Paclobutrazol	68.33 <sup>b</sup>
T <sub>12</sub> : Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl	45.00 <sup>c</sup>
T <sub>13</sub> : Control	35.00 <sup>c</sup>
<b>F- test</b>	*
<b>S.Em ±</b>	7.32
<b>CD at (5%)</b>	21.50

Figures with same alphabets did not differ significantly. \*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant. Values were compared with respective C.D values

**Table 2: Effect of Combination of Bioregulators and Growth Regulator Sprays on Panicle Length of Mango Cv. Banganpalli**

Treatment		MEAN
T <sub>1</sub> :	Gibberilic acid	21.81 <sup>b</sup>
T <sub>2</sub> :	Naphthalein Acetic Acid	29.36 <sup>ab</sup>
T <sub>3</sub> :	Paclobutrazol	30.57 <sup>a</sup>
T <sub>4</sub> :	(1-2-chloro-4pyridyl-3-phenyl urea)	20.88 <sup>b</sup>
T <sub>5</sub> :	Zinc Sulphate+Gibberilic acid	24.32 <sup>b</sup>
T <sub>6</sub> :	Zinc Sulphate+Naphthalein Acetic Acid	24.97 <sup>b</sup>
T <sub>7</sub> :	Zinc Sulphate+Paclobutrazol	26.61 <sup>ab</sup>
T <sub>8</sub> :	Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	25.52 <sup>ab</sup>
T <sub>9</sub> :	Potassium Sulphate +Gibberilic acid	24.34 <sup>b</sup>
T <sub>10</sub> :	Potassium Sulphate +Naphthalein Acetic Acid	26.96 <sup>ab</sup>
T <sub>11</sub> :	Potassium Sulphate +Paclobutrazol	25.66 <sup>ab</sup>
T <sub>12</sub> :	Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	25.46 <sup>ab</sup>
T <sub>13</sub> :	Control	19.30 <sup>b</sup>
F- test		*
S.Em ±		2.07
CD at (5%)		6.09

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values

**Table 3: Effect of Combination of Bioregulators and Growth Regulator Sprays on Panicle Breadth of Mango Cv. Banganpalli**

Treatment		Mean
T <sub>1</sub> :	Gibberilic acid	9.25 <sup>c</sup>
T <sub>2</sub> :	Naphthalein Acetic Acid	14.20 <sup>bc</sup>
T <sub>3</sub> :	Paclobutrazol	22.20 <sup>a</sup>
T <sub>4</sub> :	(1-2-chloro-4pyridyl-3-phenyl urea)	11.40 <sup>c</sup>
T <sub>5</sub> :	Zinc Sulphate+Gibberilic acid	12.98 <sup>b</sup>
T <sub>6</sub> :	Zinc Sulphate+Naphthalein Acetic Acid	17.06 <sup>b</sup>
T <sub>7</sub> :	Zinc Sulphate+Paclobutrazol	18.51 <sup>ab</sup>
T <sub>8</sub> :	Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	21.82 <sup>ab</sup>
T <sub>9</sub> :	Potassium Sulphate +Gibberilic acid	10.53 <sup>c</sup>
T <sub>10</sub> :	Potassium Sulphate +Naphthalein Acetic Acid	16.41 <sup>bc</sup>
T <sub>11</sub> :	Potassium Sulphate +Paclobutrazol	16.6 <sup>bc</sup>
T <sub>12</sub> :	Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	15.87 <sup>bc</sup>
T <sub>13</sub> :	Control	8.33 <sup>c</sup>
F- test		*
S.Em ±		1.51
CD at (5%)		4.45

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values.

**Table 4: Effect of Combinations of Bioregulators and Growth Regulator Sprays on Fruit Set Per Panicle of Mango Cv. Banganpalli**

Treatment	Mean
T <sub>1</sub> : Gibberilic acid	8.86 <sup>b</sup>
T <sub>2</sub> : Naphthalein Acetic Acid	9.06 <sup>b</sup>
T <sub>3</sub> : Pachlobutrazol	13.13 <sup>a</sup>
T <sub>4</sub> : (1-2-chloro-4pyridyl-3-phenyl urea)	9.26 <sup>b</sup>
T <sub>5</sub> : Zinc Sulphate+Gibberilic acid	10.13 <sup>ab</sup>
T <sub>6</sub> : Zinc Sulphate+Naphthalein Acetic Acid	8.20 <sup>b</sup>
T <sub>7</sub> : Zinc Sulphate+Pachlobutrazol	8.43 <sup>b</sup>
T <sub>8</sub> : Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	13.00 <sup>a</sup>
T <sub>9</sub> : Potassium Sulphate +Gibberilic acid	10.86 <sup>ab</sup>
T <sub>10</sub> : Potassium Sulphate +Naphthalein Acetic Acid	11.33 <sup>ab</sup>
T <sub>11</sub> : Potassium Sulphate +Pachlobutrazol	12.80 <sup>ab</sup>
T <sub>12</sub> : Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	8.33 <sup>b</sup>
T <sub>13</sub> : Control	8.00 <sup>b</sup>
<b>F- test</b>	*
<b>S.Em ±</b>	1.24
<b>CD at (5%)</b>	3.62

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values

**Table 5: Effect of Combination of Bioregulators and Growth Regulators Sprays on Fruit Number of Mango Cv. Banganpalli**

Treatment	Mean
T <sub>1</sub> : Gibberilic acid	113.00 <sup>c</sup>
T <sub>2</sub> : Naphthalein Acetic Acid	126.00 <sup>b</sup>
T <sub>3</sub> : Paclobutrazol	133.00 <sup>a</sup>
T <sub>4</sub> : (1-2-chloro-4pyridyl-3-phenyl urea)	131.66 <sup>ab</sup>
T <sub>5</sub> : Zinc Sulphate+Gibberilic acid	112.33 <sup>c</sup>
T <sub>6</sub> : Zinc Sulphate+Naphthalein Acetic Acid	120.66 <sup>b</sup>
T <sub>7</sub> : Zinc Sulphate+Paclobutrazol	131.33 <sup>ab</sup>
T <sub>8</sub> : Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	122.66 <sup>b</sup>
T <sub>9</sub> : Potassium Sulphate +Gibberilic acid	126.00 <sup>b</sup>
T <sub>10</sub> : Potassium Sulphate +Naphthalein Acetic Acid	123.33 <sup>b</sup>
T <sub>11</sub> : Potassium Sulphate +Paclobutrazol	130.00 <sup>ab</sup>
T <sub>12</sub> : Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	123.33 <sup>b</sup>
T <sub>13</sub> : Control	110.66 <sup>c</sup>
<b>F- test</b>	*
<b>S.Em ±</b>	2.29
<b>CD at (5%)</b>	6.68

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values



**Table 6: Effect of Combination of Bioregulators and Growth Regulator Sprays on Fruit Weight of Mango Cv. Banganpalli**

Treatment		Mean
T <sub>1</sub> :	Gibberilic acid	301.00 <sup>bc</sup>
T <sub>2</sub> :	Naphthalein Acetic Acid	315.67 <sup>a</sup>
T <sub>3</sub> :	Paclobutrazol	309.33 <sup>ab</sup>
T <sub>4</sub> :	(1-2-chloro-4pyridyl-3-phenyl urea)	299.67 <sup>bc</sup>
T <sub>5</sub> :	Zinc Sulphate+Gibberilic acid	301.67 <sup>bc</sup>
T <sub>6</sub> :	Zinc Sulphate+Naphthalein Acetic Acid	297.33 <sup>bc</sup>
T <sub>7</sub> :	Zinc Sulphate+Paclobutrazol	309.00 <sup>ab</sup>
T <sub>8</sub> :	Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	305.33 <sup>b</sup>
T <sub>9</sub> :	Potassium Sulphate +Gibberilic acid	306.33 <sup>ab</sup>
T <sub>10</sub> :	Potassium Sulphate +Naphthalein Acetic Acid	307.33 <sup>ab</sup>
T <sub>11</sub> :	Potassium Sulphate +Paclobutrazol	299.33 <sup>bc</sup>
T <sub>12</sub> :	Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	309.00 <sup>ab</sup>
T <sub>13</sub> :	Control	296.67 <sup>c</sup>
<b>F- test</b>		*
<b>S.Em ±</b>		3.10
<b>CD at (5%)</b>		9.05

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values

**Table 7: Effect of Combination of Bioregulators and Growth Regulator Sprays on Yield (Kg per Tree) of Mango of Mango Cv. Banganpalli**

Treatment		MEAN	Percentage (%) Increase in Yield Over Control
T <sub>1</sub> :	Gibberilic acid	34.01 <sup>e</sup>	3.62
T <sub>2</sub> :	Naphthalein Acetic Acid	39.7 <sup>ab</sup>	20.96
T <sub>3</sub> :	Paclobutrazol	40.6 <sup>a</sup>	23.70
T <sub>4</sub> :	(1-2-chloro-4pyridyl-3-phenyl urea)	39.45 <sup>b</sup>	20.20
T <sub>5</sub> :	Zinc Sulphate+Gibberilic acid	34.74 <sup>e</sup>	5.85
T <sub>6</sub> :	Zinc Sulphate+Naphthalein Acetic Acid	35.87 <sup>d</sup>	9.35
T <sub>7</sub> :	Zinc Sulphate+Paclobutrazol	39.61 <sup>ab</sup>	20.68
T <sub>8</sub> :	Zinc Sulphate+1-2-chloro-4pyridyl-3-phenyl urea	37.9 <sup>c</sup>	15.48
T <sub>9</sub> :	Potassium Sulphate +Gibberilic acid	38.59 <sup>bc</sup>	17.60
T <sub>10</sub> :	Potassium Sulphate +Naphthalein Acetic Acid	37.87 <sup>c</sup>	15.39
T <sub>11</sub> :	Potassium Sulphate +Paclobutrazol	38.91 <sup>bc</sup>	18.55
T <sub>12</sub> :	Potassium Sulphate +1-2-chloro-4pyridyl-3-phenyl urea)	38.11 <sup>c</sup>	16.12
T <sub>13</sub> :	Control	32.82 <sup>f</sup>	
<b>F- test</b>		*	
<b>S.Em ±</b>		0.38	
<b>CD at (5%)</b>		1.11	

Figures with same alphabets did not differ significantly.

\*\* Significant at (p= 0.01 LOS), \*Significant at (p= 0.05 LOS), NS- Non Significant.

Values were compared with respective C.D values

## CONCLUSIONS

Among the bioregulators and growth regulator sprays, highest yield was recorded in paclobutrazol (23.7 % over control) alone applied trees compare to control. Naphthalene acetic acid (NAA) spray alone has increased the yield up to 20.96 % over control was effective in improving the yield of mango. Zinc sulphate+ paclobutrazol application has increased the yield up to 20.68 % over control and (1-2-chloro-4pyridyl-3-phenylurea) application has increased the yield up to 20.20% over control. Hence, paclobutrazol and NAA alone were equally effective in improving the yield when compared to bioregulators, growth regulators sprayed in other combinations.

## REFERENCES

1. Abdollahi, M. Eshghi, S, and Tafazzoli, E. 2010. Interaction of paclobutrazol, boron and zinc on vegetative growth, yield and fruit quality of strawberry (*Fragaria x Ananassa* Duch. cv. Selva). *Journal of Biological & Environmental Sciences*. 4(11):67-75.
2. Anonymous. 2013. Annual Report, National Horticulture Board.
3. Banik, B.C and Sen, S.K. 1997. Effect of three levels of Zinc, iron, boron and their interactions on growth, flowering and yield of mango. *Horticultural journal*. 10(1):23-29.
4. Bhagwan, A, Vanajalatha, K, Prabhakar Reddy, I, Sarkar, S.K and Girwani, A 2011. Standardization of Dosage and Time of Soil Application of Cultar on Flowering and Yield of Mango cv. Banganpalli. Presented in Global Conference on Augmenting Production and Utilization of Mango: Biotic and Abiotic Stresses held on 21-24<sup>th</sup> June at Luck now.
5. Bleiholder, H., van den Boom, T., Langeluddeke, P. and Stauss, R. 1989. Einheitliche Codierung der phänologischen Stadien bei Kultur- und Schadpflanzen – Gesunde Pflanzen. 41, 381–384.
6. Burondkar, M.M, Gunjate, R.T Magdum, M.B and Goverker M.A. 2000. Rejuvenation of old and overcrowded Alphonso mango orchard with pruning and use of paclobutrazol. *Acta Horticulturae*. 509(2): 681-686.
7. Burondkar, M.M and Gunjate, R.J. 1991. Regulation of shoot growth and flowering habit in Alphonso mango with Paclobutrazol. *Acta Horticulturae*. 291: 79-84.
8. Cardenas, K and Rojas, E. 2003. Effect of paclobutrazol and nitrates of potassium and calcium on the development of the mango 'Tommy Atkins'. *Bioagro*.15: 2: 83 -90.
9. Davenport 2007. Reproductive physiology of mango. *Braz. J. Plant Physiol*. 19(4):363 376.
10. Desai, M.M and Chundawat, B.S. 1994. Regulation of flowering in mango by paclobutrazol. *Indian Journal of Horticulture*. 51(1): 9-15.
11. GoGuey, T. 1990. The effects of repeated application of cultar (Paclobutrazol) to *Mangifera indica* L. Var. Valencia. *Fruits (Paris)* 45(6): 599-607.
12. Hack, H., Bleiholder, H., Buhr, L., Meier, U., Schnock-Frick, U., Weber, E. and Witzemberger. A. 1992. Einheitliche Codierung der phänologischen Entwicklungsstadien mono- und dikotyler Pflanzen – Erweiterte BBCH-Skala, Allgemein. *Nachrichtenbl. Deut. Pflanzenschutzd*. 44, 265–270.
13. Hernandez Delgado, P. M., Aranguren, M., Reig, C., Fernandez Galvan, D., Mesejo, C., Martinez Fuentes, A., Galan Sauco, V. and Agusi, M. 2011. Phenological growth stages of mango (*Mangifera indica* L.) according to the BBCH scale. *Scientia Horticulturae*. 130: 536–540.

14. Iyer, C.P.A and Kurian, R.M. 1992. Tree size controls in mango (*Mangifera indica* L.) some considerations. *Acta Horticulturae* 321: 425-436.
15. \*Kachru, R.B, Singh, R.N and Chacko, E.K. 1971. Inhibition of flowering in mango by gibberellic acid. *Horticulture Science*. 6:140-141.
16. Kumar Raj, M, Reddy, Y.N, Chandrasekhar, R, and Srihari, D 2005. Effect of foliar application of chemicals and plant growth regulators on flowering of unpruned mango trees of cv. Baneshan. *Journal of Research ANGRAU*. 33: 2, 6-11.
17. Kumbhar, A.R, Gunjate, R.T and Amim, S.M 2007. Comparision of cultar and austar as source of paclobutrazol for flowering and fruiting in Kesar in Mango. *Acta Hort*. 820, *Proc. 8<sup>th</sup> Int. Mango symposium*
18. Lolaei, A, Rezaei, M. A, Raad, M. K, Kaviani, B. 2012. Effect of paclobutrazol and sulfate zinc on vegetative growth, yield and fruit quality of strawberry (*Fragaria x ananassa*) *Annals of Biological Research*; 3(10):4657-4662.
19. Nav Prem Singh, 2002. Effect of chemicals and plant bioregulators on the promotion of flowering and fruiting in mango cv. Dusehari. Ph.D Thesis submitted to the Punjab Agricultural University, Ludhiana.
20. Negi, S.S, Singh, A.K, Narayan Singh and Hariom Sah 2011. Effect of nutrients on flowering and fruiting of mango cv. Dashehari. *Haryana Journal of Horticultural Sciences*; 40(1/2):69-71.
21. Notodimedjo, S. 2000. Effect of GA<sub>3</sub>, NAA and CPPU on fruit retention, yield and quality of mango (cv.Arumanis) in EastJava. *Acta Horticulturae*. (509):587-600.
22. Orwintinee, Chusri Naoko Koza, Tatsushi Ogata, Hirokazu Higuchi and Yoshimi Yonemoto 2008. Application of paclobutrazol for flowering and fruit production of Irwin mango (*Mangifera indica* L.) in Okinawa. *Trop. Agr.Develop*. 52(3): 69-73.
23. Patil, H.K and Talathi, J.M. 1999. Economic viability of paclobutrazol application in mango cv. Alphonso. *South Indian Horticulturae* 47(1-6): 145-146.
24. \*Quinlin, J.D and Richardson, P.J. 1984. Effect of Paclobutrazol on apple shoots growth. *Acta Horticulture*.146:105-110.
25. Ram, S and Tripathi, P.C. 1993. Effect of cultar on flowering and fruiting in high density Dashehari mango trees. *Indian Journal of Horticulture*. 50(4): 292-295.
26. Rao, M.M, Srihari, D, Patil, V.S, and Madalageri, M.B 1997. Further studies on chemical induction of flowering directly on fruited shoots in off phase Alphonso mango trees. *Karnataka J. Sci*. 10(2): 598-601.
27. Rawash, M, El-Nas, N, El-Mosry, H and Ebbed, S 1998a. Effect of post bloom spray with some chemical substances on yield and fruit quality of Taimour mango trees. *Egyptian Journal of Horticulture*. 25(1): 71-81.
28. Razzaq, K, Khan, A.S, Malik, A.U, Shahid, M and Ullah, S. 2013. Foliar application of zinc influences the leaf mineral status, vegetative and reproductive growth, yield and fruit quality of 'Kinnow' mandarin. *Journal of Plant Nutrition*. 36(10):1479-1495. 40.
29. Shailender rajan, Divya Tiwari, Singh, V.K., Reddy, Y.T.N., Upreti, K.K, Burondkhar, M.M., Bhagwan, A., Kennedy R. and Pooja Saxena. 2011. Application of extended BBCH scale for phenological studies in mango (*Mangifera indica* L). *Journal of Applied Horticulture* 13(2) July- September
30. Shinde, A.K, Burondkar, M.M, Bhingrade, R.T, Waghmare, G.M, Rangwala, A.D and Wagh, R.G 2001. Heat unit requirement for fruit maturity in mango varieties. *Indian Journal of Plant Physiology*. 6(2): 194-196.
31. Shinde, A.K, Dalvi, M, Godse, S, Patil, B and Pujari, K 2002. Evaluation of chemical and growth regulatory for post harvest treatment of fruits in Alphonso mango. VII International Mango Symposium, Recife, Brazil, p.77.

32. Singh, A.K and Rajput C.B.S 1990. Effect of GA, BA and Calcium on vegetative growth and flowering in mango (*Mangifera indica* L.). *Research and Development Reporter* 7(1-2): 1-11.
33. Singh, N.P, Malhi, C.S, Sharma, R.C. 2005. Effect of plant bio regulators (PBRs) on flowering, fruit yield and quality in mango cv. Dashehari. *Horticultural Journal*. 18: 1, 10-12.
34. Singh, Z and Dhillon, B.S. 1992. Effect of paclobutrazol on floral malformation, yield and quality of mango (*Mangifera indica* L.). *Acta Horticulturae* 296: 51-54.
35. Vijayalaxmi, D and Srinivasan, P.S. 2000. Altering the enzyme activities for enhancing yield in 'Off' year mango (cv. Alphonso) through chemicals and growth regulators. *Orissa Journal of Horticulture*. 28(2): 1-7.
36. Vijayalakshmi, D and P.S Srinivasan. 1998. Induction of flowering in off year mango cv. Alphonso as influenced by chemicals and growth regulators. *Ann. Plant Physiol*. 12(2): 93-97.
37. Wahdan, M.T, Habib, S.E, Bassal, M.A and Qaoud E.M 2011. Effect of some chemicals on growth, fruiting, yield and fruit quality of "Succary Abiad" mango cv. *Journal of American Science*. 7(2): 651-658.
38. Winston, E.C 1992. Evaluation of paclobutrazol on growth, flowering and yield of mango cv. Kensington pride. *Australian Journal of Experimental Agriculture*. 32(1):97-104.
39. Zora Singh, Singh, Z, Muller, W, Polency F, Verheyden, C and Webster, A.D 2000. Effect of (2 RS, RS) paclobutrazol on tree vigour, flowering, fruit set and yield in mango. *Proceeding of International Conference on Integrated fruit production, Leuven, Belgium 27 July-1 August*. *Acta Horticulture* 525:459-462(Abst.).
40. Zora Singh and Dhillon, B.S. 1992. Effect of paclobutrazol on floral malformation, yield and quality of mango (*Mangifera indica* L.). *Acta Horticulture* 296: 51-53.